

超高磁場MRI診断・病態研究部門

氏名	所属	職名	取得学位	専門分野	主な論文・著作・業績
佐々木 真理	超高磁場MRI診断・病態研究部門	教授	博士（医学）	放射線科学、神経放射線診断学、磁気共鳴医学	<p>①Sasaki M, Kudo K, Honjo K, Hu JQ, Wang HB, Shintaku K: Prediction of infarct volume and neurologic outcome by using automated multiparametric perfusion-weighted magnetic resonance imaging in a primate model of permanent middle cerebral artery occlusion. <i>J Cereb Blood Flow Metab</i> 31:448–456 (2011)</p> <p>②Hirano T, Sasaki M, Mori E, Minematsu K, Nakagawara J, Yamaguchi T: Residual Vessel Length on Magnetic Resonance Angiography Identifies Poor Responders to Alteplase in Acute Middle Cerebral Artery Occlusion Patients: Exploratory Analysis of the Japan Alteplase Clinical Trial II. <i>Stroke</i> 41:2828–2833 (2010)</p> <p>③Sasaki M, Kudo K, Ogasawara K, Fujiwara S: Tracer delay-insensitive algorithm can improve reliability of CT perfusion imaging for cerebrovascular steno-occlusive disease: comparison with quantitative single-photon emission CT. <i>AJR Am J Neuroradiol</i> 30:188–193 (2009)</p> <p>④Sasaki M, Yamada K, Watanabe Y, Matsui M, Ida M, Fujiwara S, Shibata E: Variability in absolute apparent diffusion coefficient values across different platforms may be substantial: a multivendor, multi-institutional comparison study. <i>Radiology</i> 249:624–630 (2008)</p> <p>⑤Shibata E, Sasaki M, Tohyama K, Otsuka K, Endoh J, Terayama Y, Sakai A: Use of neuromelanin-sensitive MRI to distinguish schizophrenic and depressive patients and healthy individuals based on signal alterations in the substantia nigra and locus ceruleus. <i>Biol Psychiatry</i> 64:401–406 (2008)</p>
山下 典生	超高磁場MRI診断・病態研究部門	助教	博士（医学）	脳計測科学 医用システム 生命・健康・医療情報学	<p>①Yamashita F, Sasaki M, Fukumoto K, Otsuka K, Uwano I, Kameda H, Endoh J, Sakai A: Detection of changes in the ventral tegmental area of patients with schizophrenia using neuromelanin-sensitive MRI. <i>Neuroreport</i> 27(5):289–294 (2016)</p> <p>②Okada N, Fukunaga M, Yamashita F, Koshyama D, Yamamori H, Ohi K, Yasuda Y, Fujimoto M, Watanabe Y, Yahata N, Nemoto K, Hibar DP, van Erp TG, Fujino H, Isobe M, Isomura S, Natsubori T, Narita H, Hashimoto N, Miyata J, Koike S, Takahashi T, Yamasue H, Matsuo K, Onitsuka T, Iidaka T, Kawasaki Y, Yoshimura R, Watanabe Y, Suzuki M, Turner JA, Takeda M, Thompson PM, Ozaki N, Kasai K, Hashimoto R: Abnormal asymmetries in subcortical brain volume in schizophrenia. <i>Mol Psychiatry</i> doi: 10.1038/mp.2015.209. (2016)</p> <p>③Yamashita F, Sasaki M, Saito M, Mori E, Kawaguchi A, Kudo K, Natori T, Uwano I, Ito K, Saito K: Voxel-based morphometry of disproportionate cerebrospinal fluid space distribution for the differential diagnosis of idiopathic normal pressure hydrocephalus. <i>J Neuroimaging</i> 24(4):359–365 (2014)</p> <p>④Maikusa N, Yamashita F, Tanaka K, Abe O, Kawaguchi A, Kabasawa H, Chiba S, Kasahara A, Kobayashi N, Yuasa T, Sato N, Matsuda H and Iwatsubo T: Improved volumetric measurement of brain structure with a distortion correction procedure using an ADNI phantom. <i>Med Phys</i> 40(6):062303 (2013)</p> <p>⑤Yamashita F, Sasaki M, Takahashi S, Matsuda H, Kudo K, Narumi S, Terayama Y, Asada T: Detection of changes in cerebrospinal fluid space in idiopathic normal pressure hydrocephalus using voxel-based morphometry. <i>Neuroradiology</i>. 52:381–6 (2010)</p>

超高磁場MRI診断・病態研究部門

氏名	所属	職名	取得学位	専門分野	主な論文・著作・業績
樋口 さとみ	超高磁場MRI診断・病態研究部門	助教	博士（理学）	脳計測科学、計算論的神経科学	<p>①Di Dio C, Di Cesare G, Higuchi S, Roberts N, Vogt S, Rizzolatti G: The neural correlates of velocity processing during the observation of a biological effector in the parietal and premotor cortex. <i>Neuroimage</i>. 64:425–436 (2013)</p> <p>②Higuchi S, Holle H, Roberts N, Eickhoff SB, Vogt S: Imitation and observational learning of hand actions: prefrontal involvement and connectivity. <i>Neuroimage</i> 16: 1668–1683 (2012)</p> <p>③Higuchi S, Chaminade T, Imamizu H, Kawato M: Shared neural correlates for language and tool use in Broca's area. <i>Neuroreport</i>. 20:1376–1381 (2009)</p> <p>④Higuchi S, Imamizu H, Kawato M: Cerebellar activity evoked by common tool-use execution and imagery tasks: an fMRI study. <i>Cortex</i>. 43:350–358 (2007)</p> <p>⑤Imamizu H, Higuchi S, Toda A, Kawato M: Reorganization of brain activity for multiple internal models after short but intensive training. <i>Cortex</i>. 43:338–349 (2007)</p>
上野 育子	超高磁場MRI診断・病態研究部門	助教	博士（ソフトウェア情報学） 博士（医学）	画像情報処理、情報学基礎理論	<p>①Uwano I, Sasaki M, Kudo K, Boutelier T, Kameda H, Mori F, Yamashita F: Tmax Determined Using a Bayesian Estimation Deconvolution Algorithm Applied to Bolus Tracking Perfusion Imaging: A Digital Phantom Validation Study. <i>Magn Reson Med Sci.</i> (Epub 2016)</p> <p>②Uwano I, Metoki T, Sendai F, Yoshida R, Kudo K, Yamashita F, Higuchi S, Ito K, Harada T, Goodwin J, Ogawa A, Sasaki M: Assessment of Sensations Experienced by Subjects during MR Imaging Examination at 7T. <i>Magn Reson Med Sci.</i> 14(1):35–41 (2015)</p> <p>③Uwano I, Kudo K, Yamashita F, Goodwin J, Higuchi S, Ito K, Harada T, Ogawa A, Sasaki M: Intensity inhomogeneity correction for magnetic resonance imaging of human brain at 7T. <i>Med Phys</i> 41(2):022302 (2014)</p> <p>④Uwano I, Sasaki M, Kudo K, Fujiwara S, Yamaguchi M, Saito A, Ogasawara K, Ogawa A: Diffusion anisotropy color-coded map of cerebral white matter: quantitative comparison between orthogonal anisotropic diffusion-weighted imaging and diffusion tensor imaging. <i>J Neuroimaging</i> 23(2):197–201 (2013)</p> <p>⑤Uwano I, Kudo K, Sasaki M, Christensen S, Ostergaard L, Ogasawara K, Ogawa A: CT and MR perfusion can discriminate severe cerebral hypoperfusion from perfusion absence: Evaluation of different commercial software packages by using digital phantoms. <i>Neuroradiology</i> 54(5):467–474 (2012)</p>

超高磁場MRI診断・病態研究部門

氏名	所属	職名	取得学位	専門分野	主な論文・著作・業績
伊藤 賢司	超高磁場MRI診断・病態研究部門	助教（任期付）	博士（医学）	放射線技術学、脳拡散磁気共鳴画像、医用画像工学	<p>①Ito K, Kudo M, Sasaki M, Saito A, Yamashita F, Harada T, Yokosawa S, Uwano I, Kameda H, Terayama Y. Detection of changes in the periaqueductal gray matter of patients with episodic migraine using quantitative diffusion kurtosis imaging: preliminary findings. <i>Neuroradiology.</i> 58:115-20 (2016)</p> <p>②Ito K, Sasaki M, Ohtsuka C, Yokosawa S, Harada T, Uwano I, Yamashita F, Higuchi S, Terayama Y. Differentiation among parkinsonisms using quantitative diffusion kurtosis imaging. <i>Neuroreport.</i> 26:267-72 (2015)</p> <p>③Ito K, Sasaki M, Takahashi J, Uwano I, Yamashita F, Higuchi S, Goodwin J, Harada T, Kudo K, Terayama Y. Detection of early changes in the parahippocampal and posterior cingulum bundles during mild cognitive impairment by using high-resolution multi-parametric diffusion tensor imaging. <i>Psychiatry Res.</i> 231:346-52 (2015)</p> <p>④Ito K, Sasaki M, Kobayashi M, Ogasawara K, Nishihara T, Takahashi T, Natori T, Uwano I, Yamashita F, Kudo K. Noninvasive evaluation of collateral blood flow through circle of Willis in cervical carotid stenosis using selective magnetic resonance angiography. <i>J Stroke Cerebrovasc.</i> 23:1019-23 (2014)</p> <p>⑤Ito K, Masutani Y, Kamagata K, Yasmin H, Suzuki Y, Ino K, Aoki S, Kunimatsu A, Ohtomo K. Automatic extraction of the cingulum bundle in diffusion tensor tract-specific analysis: Feasibility study in Parkinson's disease with and without dementia. <i>Magn Reson Med Sci.</i> 12:201-13 (2013)</p>